

Parity and Large Gauge Invariance in Thermal QED₃

D. Seminara

Department of Physics, Brandeis University, Waltham, MA 02454, USA

We settle the “apparent” paradox present in thermal QED₃ that the perturbative series is not invariant, as manifested by the temperature dependence of the induced Chern-Simons term, by showing that large (unlike small) transformations and hence their Ward identities, are not perturbative order-preserving. Instead the thermal effective gauge field actions induced by charged fermions in QED₃ can be made invariant under both small and large gauge transformations by suitable regularization of the Dirac operator determinant, at the usual price of parity anomalies. Our result is illustrated by a concrete example.

Three-dimensional gauge theories are of physical interest in the condensed matter context¹, but display special features requiring understanding different from their four dimensional counterparts. In particular, we will be concerned with the complex of problems associated with the presence of Chern-Simons (CS) terms², the necessary quantization of their coefficients^{2,3} in the action stemming from the possibility of making homotopically nontrivial “large” gauge transformations, and the effect of quantum loop corrections on this sector⁴. While large transformations are always relevant in the nonabelian case, they also come into play in the physically most interesting case of QED₃ at finite temperatures where the compactified euclidean time/temperature provides a nontrivial, S^1 , geometry. These exotic features have been the subject of a large literature⁵, as they seemingly lead to a paradox: on the one hand, large gauge invariance appears to require quantization of the CS term’s coefficient; on the other, matter loop contributions to the effective gauge field action at finite temperatures yield a perturbative expansion in which it acquires temperature-dependent, hence non-quantized, coefficients that seems to signal a gauge anomaly. This is particularly puzzling since both the matter action and the process of integrating out its excitations should be intrinsically gauge invariant. Here, summarizing our previous results^{6,7}, we will show that the effective action is indeed invariant under both small and large transformations using classic results on elliptic operators⁸ that allow a clear definition of the Dirac operator’s functional determinant by means of ζ -function regularization. Instead, we will see that it is the perturbative expansion that is non-invariant because large transformations necessarily introduce non-analytic dependence on the charge so that expansion in e^2 and large gauge invariance are mutually incompatible: the induced Chern-Simons term’s non-invariance is precisely compensated by further non-local contributions in the effective ac-

tion. We will also note the necessary clash between gauge invariance and parity conservation, similar to that in the familiar axial anomaly in even dimensions. All these features are illustrated by explicit consideration of a non-trivial configuration that enables us to “parametrize” the CS aspects.

Let us begin with the peculiar properties of large gauge transformations that invalidate the usual Ward identity consistency. Restoring explicit dependence on e , we have $A_\mu \rightarrow A_\mu + e^{-1} \partial_\mu f$. Normally, we can merely redefine $\tilde{f} = e^{-1} f$. This is also true at finite temperature for the small gauge transformations since f is only required to be periodic in Euclidean time $\beta = (\kappa T)^{-1}$. Thus a perturbative expansion will be small gauge invariant order by order. But for large ones, the periodicity condition becomes $f(0, \mathbf{r}) = f(\beta, \mathbf{r}) + 2\pi i n$, with $n \in \mathbb{Z}$, and a rescaling will merely hide the e^{-1} factor in the boundary conditions. This intrinsic dependence means that only the *full* effective action, but not its individual expansion terms (including CS parts !) remains invariant. We are therefore driven to a careful treatment of the induced effective action $\Gamma[A]$ resulting from integrating out the charged matter, for us massive fermions, according to the usual relation $\exp(-\Gamma[A]) = \det(i\not{D} + im)$ where D_μ is the $U(1)$ covariant derivative. The extension to N flavors and to the non-abelian case will be seen to be straightforward. Our 3-space has $S^1(\text{time}) \times \Sigma$ topology, Σ being a compact Riemann 2-surface. We work with a finite 2-volume in order to avoid infrared divergences associated with the continuous spectrum in an open space. Before proceeding, let us see how assuming gauge invariance constrains the form of the determinant. [To avoid irrelevant spatial homotopies, we shall here take Σ to be the sphere.] Because of the existence of the non-trivial S^1 cycle we can construct (besides $F_{\mu\nu}$) the gauge invariant holonomy $\Omega(\mathbf{r}) \equiv \exp\left(i \int_0^\beta A_0(t', \mathbf{r}) dt'\right)$. Ω is not a completely independent variable, as part of the information carried by it is already present in $F_{\mu\nu}$: it satisfies the constraint $\nabla \Omega = i \Omega \int_0^\beta \mathbf{E}(t', \mathbf{r}) dt'$, implying that Ω has the form $\Omega = \exp(2\pi i a) \Omega_0(\mathbf{E})$, where $\Omega_0(\mathbf{E})$ is a nonlocal functional depending only on $\mathbf{E}$ and on the geometry of S^2 . The new information is encoded entirely in the constant a , the flat connection. [For example, the non trivial behavior of A_0 under large gauge transformation is inherited by a : $a \rightarrow a + 1$.] Therefore the determinant can be considered as a function(al) of $F_{\mu\nu}$ and a alone. Large gauge invariance implies the separate Ward identity $e^{-\Gamma(a+1, F)} = e^{-\Gamma(a, F)}$, namely periodicity. Then Fourier-expanding and factorizing out the parity anomaly contribution, we obtain

$$e^{-\Gamma(F, a)} = e^{i S_{CS}} \sum_{k=0}^{\infty} \Gamma_k^{(1)}(F_{\mu\nu}) e^{\pi(2k - \Phi(F))a} + \Gamma_k^{(2)}(F_{\mu\nu}) e^{-\pi(2k - \Phi(F))}, \quad (1)$$

where $\Phi(F)$ is the electromagnetic flux through S^2 and S_{CS} is the abelian CS action. [The correct form for the abelian CS action, when A carries a nontrivial flux, hides some subtleties that have been reviewed in^{7,9}.] As we shall see, the structure exhibited in (1) will be explicitly realized in our example.

We now return to the definition of the effective action. Within our framework, the Dirac operator is a well-defined elliptic operator⁸ whose determinant can be rigorously specified. The ζ -function regularization¹⁰ defines the formal product of all the eigenvalues λ_n as

$$\det i(\not{D} + m) = \Pi \lambda_n \equiv \exp(-\zeta'(0)), \quad \zeta(s) \equiv \sum (\lambda_n)^{-s} \quad (2)$$

with implicit repetition over degenerate eigenvalues. For $s > 3$ in $D = 3$ ⁸, the above series converges and its analytic extension defines a meromorphic function with only simple poles. It is regular at $s = 0$, thereby assuring the meaningfulness of (2). A careful definition of λ_n^{-s} is required to avoid ambiguities. We take it to be $\exp(-s \log \lambda_n)$ where the cut is chosen to be over the positive real axis, $0 \leq \arg \lambda_n < 2\pi$, enabling us to rewrite $\zeta(s)$ in the more convenient form

$$\zeta(s) = \sum_{\text{Re } \lambda_n > 0} (\lambda_n)^{-s} + \exp(-i\pi s) \sum_{\text{Re } \lambda_n < 0} (-\lambda_n)^{-s}. \quad (3)$$

Changing the cut only alters the determinant if it intersects the line $\text{Im} z = m$, in which case the only relevant difference is the sign of the exponential in (3). This alternative choice does not affect gauge invariance, but does change the sign of the parity anomaly terms in $\Gamma[A]$. Once the determinant of the Dirac operator has been regularized, its full gauge invariance reduces to that of its eigenvalue spectrum. But small transformations do not affect the λ_n at all, while the large ones merely permute them, as in usual illustrations of index theorems⁸; every well-defined symmetric function of the spectrum, such as $\zeta(s)$ and hence $\Gamma[A]$, is unchanged.

The price paid for preserving gauge invariance is (as usual !) an intrinsic parity anomaly, *i.e.*, one present even in the limit when the explicitly parity violating fermion mass term is absent. [That the parity can be sacrificed for gauge was effectively noted in¹¹.] Under P , $\lambda_n \rightarrow -\lambda_n^*$ so that $\zeta^P(s) \neq \zeta(s)$. It is easy to express the parity violating part $\Gamma^{(PV)}[A] = 1/2(\zeta'(0) - \zeta'^P(0))$ explicitly in terms of the η -function in this limit. Here, $\zeta^{PV} \equiv \zeta^P - \zeta$ is

$$\zeta^{PV}(s) = (1 - e^{-i\pi s}) \left(\sum_{\lambda_n > 0} (\lambda_n)^{-s} - \sum_{\lambda_n < 0} (-\lambda_n)^{-s} \right) \equiv (1 - e^{-i\pi s}) \eta(s), \quad (4)$$

so that $\Gamma^{(PV)}[A] = i\pi/2\eta(0)$. At $m = 0$, the continuous part of $\eta(0)$ is given in closed form by the CS action⁸; being local means it can be removed by a different choice of regularization. For $m \neq 0$ an expansion in powers of the mass can be presented

$$\Gamma^{(PV)}(A) = \frac{1}{2} \frac{d}{ds} (\zeta^{PV}(s)) \Big|_{s=0} = i\frac{\pi}{2}\eta(0) - i \sum_{k=0}^{\infty} (-1)^k \frac{m^{(2k+1)}}{2k+1} \eta(2k+1), \quad (5)$$

while the analogous expansion for the parity-conserving part involves even powers of the mass. A detailed discussion of mass expansions and the relevant references can be found in⁷.

A simple, but realistic, $(2+1)$ example^{6,7}, that illustrates the previous analysis, is the $U(1)$ field

$$A_\mu(t, \mathbf{r}) \equiv \left(\frac{2\pi}{\beta} a, \mathbf{A}(\mathbf{r}) \right), \quad (6)$$

where a is a flat connection along S^1 . $\mathbf{A}$ lives on Σ , with non-vanishing, necessarily integer, flux $\Phi(F) = n$. We concentrate on large transformations $a \rightarrow a+1$, although in higher genus Σ one could also have large transformations affecting $\mathbf{A}$. [There is an intriguing $(0+1)$ antecedent of this example^{12,7,13}.] Because of the time independence, we have a tractable eigenvalue equation for λ_n . After some work, it follows that the effective action factorizes into contributions: one depending only on the holonomy $\exp(2\pi ia)$ and the other on the value of $\mathbf{A}$ on Σ ,

$$e^{-\Gamma(A)} = [e^{-\beta m + 2\pi ia} + 1]^{\nu_+} [e^{-\beta m - 2\pi ia} + 1]^{\nu_-} \prod_{\mu_k} \left(1 + e^{-\beta \sqrt{\mu_k^2 + m^2} + 2\pi ia} \right)^2 e^{\frac{2\pi}{\beta} \zeta_{\frac{\beta^2}{4\pi^2}}(\hat{\mathcal{D}}^2 + m^2) \left((-1/2) - (\nu_+ + \nu_-) m \beta \right)}. \quad (7)$$

Here $\hat{\mathcal{D}}$ is the reduced Dirac operator on Σ , μ_k its nonvanishing eigenvalues. The number of positive/negative chiral zero-modes $v_\pm$ of $\hat{\mathcal{D}}$ is represented by $\nu_\pm$, with the conventions $(\gamma_5 \mp 1)v_\pm = 0$, and the (parity odd) flux is just $\nu_- - \nu_+$. That the infinite product in (8) is convergent follows from the fact that⁸ $\mu_k \simeq c\sqrt{|k|}$. The invariance of (7) under $a \rightarrow a+1$ is manifest and its structure is consistent with (1). It is clear that a perturbative (i.e., in power of a) expansion of (7) loses periodicity in a and hence does not see large invariance order by order. For example the Chern-Simons term ($S_{CS} = \pi a n$) has a coefficient $1 - \tanh\left(\frac{\beta m}{2}\right)$. The usually quoted coefficient omits the 1 that represents the intrinsic parity-anomaly price of our gauge-invariant

regularization and hence persists at $m = 0$. There is actually an ambiguity in its sign (reflecting the choice of cut in (3)), also present in other regularizations, for example through the factor $\lim_{M \rightarrow \pm\infty} \text{sign}(M)$ in Pauli-Villars.

In conclusion, we have shown that the apparent large gauge anomalies resulting from a perturbative expansion of the full effective action are due to the more complicated (order-violating) nature of the Ward identities when a non-trivial homotopy is present, the action itself being fully gauge invariant with suitable regularization, one that necessarily entails parity anomalies.

It is a pleasure to thank my collaborators S. Deser and L. Griguolo. This work was supported by NSF grant PHY-9315811.

References

1. J. D. Lykken, J. Sonnenschein and N. Weiss, *Int. Jour. Mod. Phys. A* **6**, 5155 (1991).
2. S. Deser, R. Jackiw and S. Templeton, *Phys. Rev. Lett.* **48**, 975 (1982); *Ann. Phys. (N.Y.)* **140**, 372 (1982).
3. S. Elitzur, G. Moore, A. Schwimmer and N. Seiberg, *Nucl. Phys. B* **326**, 108 (1989); V. P. Nair and M. Bos, *Phys. Lett. B* **223**, 61 (1989); G. Dunne, R. Jackiw and C. Trugenberger, *Ann. Phys.* **194**, 197 (1989).
4. A. Niemi and G. W. Semenoff, *Phys. Rev. Lett.* **51**, 2077 (1983); R. Pisarski and S. Rao *Phys. Rev. D* **32**, 2081 (1985); R. Pisarski, *ibid.* **35**, 664 (1987).
5. K. Babu, A. Das and P. Panigrahi, *Phys. Rev. D* **36**, 3725 (1987); I. Aitchison and J. Zuk, *Ann. Phys.* **242**, 77 (1995); N. Bralić, C. Fosco and F. Schaposnik, *Phys. Lett. B* **383**, 199 (1996); D. Cabra, E. Fradkin, G. Rossini, and F. Schaposnik, *Phys. Lett. B* **383**, 434 (1996).
6. S. Deser, L. Griguolo and D. Seminara, *Phys. Rev. Lett.* **79**, 1976 (1997).
7. S. Deser, L. Griguolo and D. Seminara, *Phys. Rev. D* **57**, 7444 (1998).
8. P. B. Gilkey, “*Invariance theory, the heat-equation and the Atiyah-Singer index theorem*”, CRC Press, Boca Raton, 1995.
9. S. Deser, L. Griguolo and D. Seminara, *Comm. Math. Phys.* **197**, 443 (1998).
10. S. W. Hawking, *Comm. Math. Phys.* **55**, 133 (1975).
11. A. N. Redlich, *Phys. Rev. Lett.* **52**, 18 (1984); *Phys. Rev. D* **29**, 2366 (1984).
12. G. Dunne, K. Lee and C. Lu, *Phys. Rev. Lett.* **78**, 3434 (1997).
13. R. Jackiw and S. Y. Pi, *Phys. Lett. B* **423**, 364 (1998).